

Comment on “Fragmented Condensate Ground State of Trapped Weakly Interacting Bosons in Two Dimensions”

Recently Liu *et al.* [1] examined the lowest state of a weakly-interacting Bose-Einstein condensate. In addition to other interesting results, using the method of the pair correlation function [2], they questioned the validity of the mean-field picture of the formation of vortices and stated that the vortices are generated at the center of the cloud. This is in apparent contradiction to the Gross-Pitaevskii approach, which predicts that the vortices successively enter the cloud from its outer parts as L/N (where N is the number of atoms in the trap and $\hbar L$ is the angular momentum of the system) increases [3,4]. We have managed to reproduce the results of Ref. [1], however a more careful analysis presented below confirms the validity of the mean-field approach.

Figures 1 and 2 show equidensity lines of

$$P(\mathbf{r}, \mathbf{r}_A) = \frac{\langle \Psi_{L,N} | \sum_{i \neq j} \delta(\mathbf{r} - \mathbf{r}_i) \delta(\mathbf{r}_A - \mathbf{r}_j) | \Psi_{L,N} \rangle}{(N-1) \langle \Psi_{L,N} | \sum_j \delta(\mathbf{r}_A - \mathbf{r}_j) | \Psi_{L,N} \rangle}, \quad (1)$$

calculated within the space of N atoms and L oscillator quanta and with the additional truncation that only considers single-particle components with no radial nodes and angular-momentum quantum numbers $m = 0, \dots, 6$ (as was also assumed in Ref. [1].) Here $|\Psi_{L,N}\rangle$ denotes the many-body wavefunction. In these figures $\mathbf{r}_A = (x, y) = (3, 0)$ in units of the oscillator length. The blue colour denotes regions of lower density, while the red colour denotes regions of higher density.

More specifically, in Fig. 1 we have taken $N = 40$ particles and $L = 28, 32, 36$, and 40 , i.e., $L/N = 0.7, 0.8, 0.9$, and 1.0 . Also Fig. 2 shows $P(\mathbf{r}, \mathbf{r}_A)$ for $L = 62, 64, 72$, and 84 , i.e., for $L/N = 1.55, 1.6, 1.8$, and 2.1 . These figures exhibit two significant differences in comparison with the conclusions of Ref. [1]: (i) their qualitative features are *not* independent of the radial position of the reference point $\mathbf{r}_A$, and in fact the important broken symmetry observed for $L/N = 0.7, 0.8$, and 0.9 is not visible, unless $\mathbf{r}_A$ is given a significantly large value, and (ii) the entry of the first vortex in the region $L/N < 1$, as well as that of the second vortex in the region $1 < L/N < 1.7$ is clearly seen to be from the outside and gradually moving inwards with increasing L/N .

Finally, Fig. 1 with $L/N = 1$ corresponds to a single-vortex-like state and is slightly asymmetric with respect to the center because of the finite number of atoms we have considered (the same is also true for the slight asymmetries in the two-fold and the three-fold symmetric states shown in Fig. 2). We have confirmed numerically that this asymmetry decreases with increasing N .

In summary, we disagree with the conclusions of Ref. [1] in the comparison of exact and mean-field description of the mode of entry of vortices into the condensate with increasing angular momentum. This does not however affect the conclusions of Ref. [1] concerning the fragmentation of a rotating condensate.

G. M. Kavoulakis and S. M. Reimann
Mathematical Physics,
Lund Institute of Technology, P.O. Box 118,
S-22100 Lund, Sweden

B. Mottelson
NORDITA,
Blegdamsvej 17, DK-2100 Copenhagen Ø, Denmark

-
- [1] X. Liu, H. Hu, L. Chang, W. Zhang, S.-Q. Li, and Y.-Z. Wang, Phys. Rev. Lett. **87**, 030404 (2001).
 - [2] G. S. Ezra and R. S. Berry, Phys. Rev. A **28**, 1974 (1983).
 - [3] D. A. Butts, and D. S. Rokhsar, Nature **397**, 327 (1999).
 - [4] G. M. Kavoulakis, B. Mottelson, and C. J. Pethick, Phys. Rev. A **62**, 63605 (2000).




FIG. 1. Equidensity lines of $P(\mathbf{r}, \mathbf{r}_A)$ for $N = 40$ and $L = 28, 32, 36$, and 40 , calculated with the reference point located at $\mathbf{r}_A = (3, 0)$.




FIG. 2. Same as Fig. 1 for $N = 40$ and $L = 62, 64, 72$, and 84 .